Equivalence between fractional exclusion statistics and Fermi liquid theory in interacting particle systems

arXiv:1303.5493v1 [cond-mat.stat-mech] 22 Mar 2013

D. V. Anghel[1] and G. A. Nemnes[1,2]

[1] *Horia Hulubei National Institute for Physics and Nuclear Engineering, P.O. Box MG-6, 077126 Măgurele, Ilfov, Romania*

[2] *University of Bucharest, Faculty of Physics, "Materials and Devices for Electronics and Optoelectronics" Research Center, P.O. Box MG-11, 077125 Măgurele-Ilfov, Romania*

PACS `05.30.-d` – Quantum statistical mechanics
PACS `05.30.Ch` – Quantum ensemble theory
PACS `05.30.Pr` – Fractional statistics systems (anyons, etc.)

Abstract – We develop a method based on fractional exclusion statistics (FES) to describe s-dimensional systems of interacting particles. The particles – which are described in the quasi-classical approximation – are in an external potential and experience a generic particle-particle interaction. We define the FES quasiparticle energies, we calculate the FES parameters of the system and we deduce the equations for the equilibrium particle populations. The FES gas is "ideal", in the sense that the quasiparticle energies do not depend on the other quasiparticle levels populations and the sum of the quasiparticle energies is equal to the total energy of the system. We prove that the FES formalism is equivalent to the standard Landau's Fermi liquid theory (FLT) approach and the FES quasiparticle populations may be calculated from the FLT populations by making the correspondence between the FES and the FLT quasiparticle energies. The FES provides a natural ideal gas description of the interacting particle gas in a generic external potential and in any number of dimensions.

Introduction. – The concept of fractional exclusion statistics (FES) was introduced by Haldane in Ref. [1] and the statistical mechanics of FES systems was formulated by several authors, employing different methods [2–7].

The FES was applied to quasiparticle excitations at the lowest Landau level in the fractional quantum Hall effect, spinon excitations in a spin-$\frac{1}{2}$ quantum antiferomagnet [1,4,8], Bose and Fermi systems described in the thermodynamic Bethe ansatz [5,9–11], excitations [1] or motifs of spins [12,13] in spin chains, elementary volumes obtained by coarse-graining in the phase-space of a system [5,14–18], interacting particles described in the mean-field approximation [19–21]. Thermodynamic properties and correlations in generic FES systems in different numbers of dimensions or interacting with external fields have also been calculated [22,23,23–27,27,28,28–31]. In Refs. [19,32] it was shown that all the systems with the same, constant density of states and diagonal, constant FES parameters (FES is manifested within each species) have the same entropy and heat capacity. This property extends the Bose-Fermi thermodynamic equivalence in 2D systems discovered long time ago [33–35], to the FES systems [36].

A stochastic method for the simulation of the time evolution of FES systems was introduced in Ref. [37] as a generalization of a similar method used for Bose and Fermi systems [38] whereas the relatively recent experimental realization of the Fermi degeneracy in cold atomic gases have renewed the interest in the theoretical investigation of non-ideal Fermi systems at low temperatures and their interpretation as ideal FES systems [27,28,39–41].

The general FES formalism was amended to include the change of the FES parameters at the change of the particle species [16,42–44]. This amendment allows the general implementation of FES as a method for the description of interacting particle systems as ideal (quasi)particle systems [17,45–47].

In this paper we apply FES to a system of particles with general particle-particle interaction, $V(|\mathbf{r}-\mathbf{r}'|)$, in an arbitrary external potential, $V_{\rm ext}(\mathbf{r})$, described in the quasi-classical approximation. We define quasiparticle energies which determine our FES parameters and using these we calculate the FES equilibrium particle distribution. We compare these results with the standard Fermi liquid theory (FLT) calculations.

The structure of the paper is as follows. In the next section we introduce our model and calculate the equilibrium particle population in the FLT approach. In the following section we implement the FES description by introducing new quasiparticle energies and the definition of species. These species are related by the FES parameters, which we calculate. Using the FES parameters and quasiparticle energies we write the equations for the equilibrium particle populations. By making the correspondence between the FLT and the FES quasiparticle energies corresponding to the same single-particle states, we show that the FES equations are satisfied by the FLT populations. This proves that the FES formalism is consistent and suitable for the description of such interacting particle systems and that the FES quasiparticle populations may be calculated using the correspondence between the quasiparticle energies.

The interacting particle system may be described as an ideal FES gas in a rather general formalism.

The model and the FLT approach. – We analyze a quasiclassical system of particles (bosons or fermions) with a generic interaction, in an external potential and s-dimensional space. We assume that the particle-particle interaction, $V(|\mathbf{r}-\mathbf{r}'|)$, depends only on the distance between the particles, whereas the external potential is $V_{\rm ext}(\mathbf{r})$. If we denote by ϵ the energies of the non-interacting particles, in the continuous limit the total energy of the system is

$$\begin{aligned} E &= \int_\Omega d^s\mathbf{r} \int_0^\infty d\epsilon\, [\epsilon + V_{\rm ext}(\mathbf{r})]\sigma(\mathbf{r},\epsilon) n(\mathbf{r},\epsilon) \\ &\quad + \frac{1}{2}\int_\Omega d^s\mathbf{r} \int_\Omega d^s\mathbf{r}' \int_0^\infty d\epsilon \int_0^\infty d\epsilon'\, \sigma(\mathbf{r},\epsilon) \\ &\quad \times \sigma(\mathbf{r}',\epsilon') n(\mathbf{r},\epsilon) n(\mathbf{r}',\epsilon') V(|\mathbf{r}-\mathbf{r}'|), \end{aligned} \tag{1}$$

where Ω is the total volume of the system and $n(\mathbf{r},\epsilon)$ is the particle population of the level ϵ in the elementary volume $d^s\mathbf{r}$. In an elementary volume, $\delta\Omega_{\mathbf{r}}$, centered at $\mathbf{r}$, the density of states (DOS) along the ϵ axis is $\delta\Omega_{\mathbf{r}}\sigma(\mathbf{r},\epsilon)$. The logarithm of the grandcanonical partition function is

$$\begin{aligned} \ln(\mathcal{Z}) &= \mp \int_\Omega d^s\mathbf{r} \int_0^\infty d\epsilon\, \sigma(\mathbf{r},\epsilon) \Big\{ [1 \mp n^{(\pm)}(\mathbf{r},\epsilon)] \\ &\quad \times \log[1 \mp n^{(\pm)}(\mathbf{r},\epsilon)] \pm n^{(\pm)}(\mathbf{r},\epsilon) \log n^{(\pm)}(\mathbf{r},\epsilon) \Big\} \\ &\quad - \beta(E - \mu N), \end{aligned} \tag{2}$$

where the upper and lower signs correspond to fermions and bosons, respectively.

The standard way to obtain the thermodynamics of the system is to maximize $\mathcal{Z}$ with respect to $n^{(\pm)}(\mathbf{r},\epsilon)$. In this way we obtain the equilibrium populations,

$$n^{(\pm)}(\mathbf{r},\epsilon) = \frac{1}{e^{\beta(\tilde\epsilon^L_{\mathbf{r}} - \mu)} \pm 1} \tag{3}$$

where

$$\tilde\epsilon^L_{\mathbf{r}} \equiv \epsilon + V_{\rm ext}(\mathbf{r}) + \int_\Omega d^s\mathbf{r}' \int_0^\infty d\epsilon' \sigma(\mathbf{r}',\epsilon') n(\mathbf{r}',\epsilon') V(|\mathbf{r}-\mathbf{r}'|) \tag{4}$$

are the Landau's quasiparticle energies.

The equilibrium distribution of particles, $n(\mathbf{r},\epsilon)$, is determined by solving self-consistently Eqs. (3) and (4). Plugging $n(\mathbf{r},\epsilon)$ back into E (1) and $\ln(\mathcal{Z})$ (2), one can calculate the total energy of the system, the grandcanonical potential and from these, the other thermodynamic quantities. We observe that the sum of the quasiparticle energies is

$$\begin{aligned} E^L &\equiv \int_\Omega d^s\mathbf{r} \int_0^\infty d\epsilon\, \tilde\epsilon^L_{\mathbf{r}}(\epsilon) \sigma(\mathbf{r},\epsilon) n(\mathbf{r},\epsilon) = E \\ &\quad + \frac{1}{2}\int_\Omega d^s\mathbf{r} \int_\Omega d^s\mathbf{r}' \int_0^\infty d\epsilon \int_0^\infty d\epsilon'\, \sigma(\mathbf{r},\epsilon) \\ &\quad \times \sigma(\mathbf{r}',\epsilon') n(\mathbf{r},\epsilon) n(\mathbf{r}',\epsilon') V(|\mathbf{r}-\mathbf{r}'|), \end{aligned} \tag{5}$$

which is different from the total energy E.

The FES formalism. – In the FES, the quasiparticle energies are defined in such a way that the total energy of the system is equal to the sum of quasiparticle energies [17,45,47,48]. For this purpose we define new quasiparticle energies,

$$\begin{aligned} \tilde\epsilon_{\mathbf{r}} &\equiv \epsilon + V_{\rm ext}(\mathbf{r}) + \int\limits_\Omega d^s\mathbf{r}' \theta[\epsilon + V_{\rm ext}(\mathbf{r}) - V_{\rm ext}(\mathbf{r}')] \\ &\quad \times \int\limits_0^{\epsilon + V_{\rm ext}(\mathbf{r}) - V_{\rm ext}(\mathbf{r}')} d\epsilon' \sigma(\mathbf{r}',\epsilon') n(\mathbf{r}',\epsilon') V(|\mathbf{r}-\mathbf{r}'|) \end{aligned} \tag{6}$$

From Eq. (6) we obtain the DOS along the $\tilde\epsilon$ axis,

$$\begin{aligned} &\tilde\sigma[\mathbf{r},\tilde\epsilon_{\mathbf{r}}(\epsilon)] = \sigma(\mathbf{r},\epsilon) \left|\frac{d\tilde\epsilon_{\mathbf{r}}}{d\epsilon}\right|^{-1} = \sigma(\mathbf{r},\epsilon) \\ &\times \Big|1 + \int_\Omega d^s\mathbf{r}'\, \theta[V_{\rm ext}(\mathbf{r}) - V_{\rm ext}(\mathbf{r}')] V(|\mathbf{r}-\mathbf{r}'|) \\ &\sigma(\mathbf{r}',\epsilon) n[\mathbf{r}',\epsilon(\tilde\epsilon_{\mathbf{r}})]\Big|^{-1}. \end{aligned} \tag{7}$$

To apply FES we split the system into imaginary elementary volumes, $\delta\Omega_{\mathbf{r}}$ (centered at $\mathbf{r}$), and in each such volume the quasiparticle energy axis, $\tilde\epsilon$, is split into elementary intervals, $\delta\tilde\epsilon$ (centered at $\tilde\epsilon$). Each such $(s+1)$-dimensional elementary volume, $\delta\Omega_{\mathbf{r}} \times \delta\tilde\epsilon_{\mathbf{r}}$, represents a FES species.

From Eqs. (6) and (7) and applying the procedure of Ref. [17, 45] we obtain the FES parameters

$$\alpha_{\delta\Omega_{\mathbf{r}}\tilde\epsilon;\delta\Omega_{\mathbf{r}'}\tilde\epsilon} = V(|\mathbf{r}-\mathbf{r}'|)\sigma[\mathbf{r},\epsilon(\tilde\epsilon)]\delta\Omega_{\mathbf{r}} \tag{8a}$$

$$\begin{aligned} \alpha_{\delta\Omega_{\mathbf{r}}\tilde\epsilon;\delta\Omega_{\mathbf{r}'}\tilde\epsilon'} &= \theta(\tilde\epsilon_{\mathbf{r}} - \tilde\epsilon'_{\mathbf{r}'}) V(|\mathbf{r}-\mathbf{r}'|) \left.\frac{d\{\ln[\sigma(\mathbf{r},\epsilon)]\}}{d\epsilon}\right|_{\epsilon(\tilde\epsilon_{\mathbf{r}})} \\ &\quad \times \sigma[\mathbf{r},\epsilon(\tilde\epsilon)]\delta\Omega_{\mathbf{r}}\delta\epsilon \equiv a_{\mathbf{r}\tilde\epsilon;\mathbf{r}'\tilde\epsilon'}\delta G(\epsilon), \end{aligned} \tag{8b}$$

$$\alpha_{\delta\Omega_{\mathbf{r}}\tilde\epsilon;\delta\Omega_{\mathbf{r}'}\tilde\epsilon'} = \theta(\tilde\epsilon_{\mathbf{r}} - \tilde\epsilon'_{\mathbf{r}'}) V(|\mathbf{r}-\mathbf{r}'|)\sigma_{\mathbf{r}}(0)\delta\Omega_{\mathbf{r}}. \tag{8c}$$

The partition function is the same as the one in Eq. (2), with $n^{(\pm)}(\mathbf{r},\epsilon) \equiv n^{(\pm)}[\mathbf{r},\tilde\epsilon_{\mathbf{r}}(\epsilon)]$, by definition [18]. Maximizing $\mathcal{Z}$ with respect to the *population along the* $\tilde\epsilon$ *axis*, we get the equations for the equilibrium particle population [18]

$$\begin{aligned}
&\beta(\mu-\tilde\epsilon_{\mathbf{r}}) + \ln\frac{[1\mp n^{(\pm)}(\mathbf{r},\tilde\epsilon_{\mathbf{r}})]}{n^{(\pm)}(\mathbf{r},\tilde\epsilon_{\mathbf{r}})} = \mp\int_\Omega d^s\mathbf{r}'\theta[\tilde\epsilon_{\mathbf{r}} - V_{\rm ext}(\mathbf{r}')] \\
&\times V(|\mathbf{r}-\mathbf{r}'|)\Big\{\sigma[\mathbf{r}',\epsilon(\tilde\epsilon_{\mathbf{r}})]\ln[1\mp n^{(\pm)}(\mathbf{r}',\tilde\epsilon_{\mathbf{r}})] \\
&+\int_{\epsilon(\tilde\epsilon_{\mathbf{r}})}^\infty d\epsilon'\,\frac{\partial\sigma(\mathbf{r}',\epsilon')}{\partial\epsilon'}\ln[1\mp n^{(\pm)}(\mathbf{r}',\epsilon')]\Big\} \\
&\mp\int_\Omega d^s\mathbf{r}'\theta[V_{\rm ext}(\mathbf{r}')-\tilde\epsilon_{\mathbf{r}}]V(|\mathbf{r}-\mathbf{r}'|) \\
&\times\Big\{\sigma(\mathbf{r}',0)\ln[1\mp n^{(\pm)}(\mathbf{r}',0)] \\
&+\int_0^\infty d\epsilon'\,\frac{\partial\sigma(\mathbf{r}',\epsilon')}{\partial\epsilon'}\ln[1\mp n^{(\pm)}(\mathbf{r}',\epsilon')]\Big\}. \qquad (9)
\end{aligned}$$

Now we can check the equivalence between the FES and the FLT descriptions of the system by substituting Eq. (3) into Eq. (9). By doing so we recover the relation between $\tilde\epsilon_{\mathbf{r}}$ and $\tilde\epsilon^L_{\mathbf{r}}$,

$$\begin{aligned}
\tilde\epsilon_{\mathbf{r}} \;\equiv\; & \tilde\epsilon^L_{\mathbf{r}} - \int_\Omega d^s\mathbf{r}'\Big\{\theta[\tilde\epsilon_{\mathbf{r}}-V_{\rm ext}(\mathbf{r}')]\int_{\epsilon(\tilde\epsilon_{\mathbf{r}})}^\infty d\epsilon'\sigma(\mathbf{r}',\epsilon') \\
&\times n(\mathbf{r}',\epsilon')V(|\mathbf{r}-\mathbf{r}'|) + \theta[V_{\rm ext}(\mathbf{r}')-\tilde\epsilon_{\mathbf{r}}] \\
&\times\int_0^\infty d\epsilon'\sigma(\mathbf{r}',\epsilon')n(\mathbf{r}',\epsilon')V(|\mathbf{r}-\mathbf{r}'|), \qquad (10)
\end{aligned}$$

which is in accordance with the definitions (4) and (6).

We illustrate our model and the relation between the FES and the FLT descriptions on a one dimensional system of fermions with repulsive Coulomb interactions – $V(|\mathbf{r}-\mathbf{r}'|) = 1/|\mathbf{r}-\mathbf{r}'|$, in the absence of external fields. The length of the system, Ω, is discretized as in Ref. [49] into $N_{\mathbf{r}}$ equal elementary segments, $\delta\mathbf{r}_\xi = 1$, where $\xi = 1,\ldots,N_{\mathbf{r}}$. In each such elementary "volume" the DOS is taken to be constant, $\delta\mathbf{r}_\xi\sigma(\mathbf{r}_\xi,\epsilon)\equiv\sigma_0$, and on the $\tilde\epsilon$ axis we define $N_{\tilde\epsilon}$ equal consecutive segments between 0 and $\tilde\epsilon_{\rm max}$, $\delta\tilde\epsilon_i \equiv \tilde\epsilon_{\rm max}/N_{\tilde\epsilon}$, where $i = 1,\ldots,N_{\tilde\epsilon}$ – we choose $\tilde\epsilon_{\rm max}$ such that $n^{(+)}(\mathbf{r},\tilde\epsilon_{\rm max}) \ll 1$ for any $\mathbf{r}$. In this way we obtain $N_{\mathbf{r}}\times N_{\tilde\epsilon}$ species of particles, $\delta\mathbf{r}_\xi\otimes\delta\tilde\epsilon_i$, identified also by a double index, (ξ,i) [49]. To avoid the singularity at the origin of the interaction potential, we consider a cut-off distance of $\delta\mathbf{r}_\xi/2$ – the choice of the cut-off does not affect the procedure.

The total number of particles in the system is $N = N_{\mathbf{r}}\epsilon_F\sigma_0$, where ϵ_F is the Fermi energy in the noninteracting system. We set the energy scale of the system by fixing $\epsilon_F = 1$ and $k_BT = 1/\beta = 1$.



Fig. 1: (Color online) Specific FES and FLT quantities for a one dimensional system with repulsive Coulomb interactions ($N_{\mathbf{r}} = 20$, $N_{\tilde\epsilon} = 50$, $\tilde\epsilon_{\rm max} = 25$): (a) the quasiparticle density of states in the FES description; (b) the position dependent energy shift, $\tilde\epsilon^L_{\xi,\rm min}$, of the Landau quasiparticle energies; (c) the FES populations; (d) the particle density, scaled with $\epsilon_F\sigma_0$ – the dashed line represents the uniform distribution of a similar non-interacting system. In the (a) and (c) subplots we plot a curve for each elementary segment, $\delta\mathbf{r}_\xi$, $\xi = 1\ldots,20$. We use the same color for symmetric segments, namely *red* for 1 and $N_{\mathbf{r}}$, *green* for 2 and $N_{\mathbf{r}}-1$, *blue* for 3 and $N_{\mathbf{r}}-2$, and black for the rest. The chemical potential, μ, is marked by vertical dashed lines.

Figure 1(a) shows the quasiparticle density of states in the FES description, $\tilde\sigma_{\xi i}\equiv\delta\mathbf{r}_\xi\tilde\sigma(\mathbf{r}_\xi,\tilde\epsilon_i)$ (Eq. 7). The density of Landau's quasiparticle states may be calculated also from Eq. (7) and we obtain $\tilde\sigma^L_{\xi i}(\tilde\epsilon^L) = \theta(\tilde\epsilon^L - \tilde\epsilon^L_{\xi,\rm min})\sigma_0$, which is different from zero only for $\tilde\epsilon^L \geq \tilde\epsilon^L_{\xi,\rm min}$. The minimum value of $\tilde\epsilon^L$ (4) is $\tilde\epsilon^L_{\xi,\rm min}\equiv\tilde\epsilon^L_{\mathbf{r}_\xi}(\epsilon=0) = \tilde\epsilon^L_{\mathbf{r}_\xi}(\epsilon)-\epsilon$. The position dependent $\tilde\epsilon^L_{\xi,\rm min}$ is plotted in Fig. 1 (b).

Figure 1(c) shows the populations, $n^{(+)}_{\xi i}\equiv n^{(+)}(\mathbf{r}_\xi,\tilde\epsilon_i)$ (9). To obtain the Landau's populations (3), we simply shift the quasiparticle energy from $\tilde\epsilon_i$ (6) to $\tilde\epsilon^L_i$ (4) and $n^{(+)}[\mathbf{r}_\xi,\tilde\epsilon_i(\tilde\epsilon^L_i)]$ becomes a Fermi distribution in $\tilde\epsilon^L_i$ for any ξ and with the same μ as for the FES distribution. The difference between the distributions $n^{(+)}(\mathbf{r}_\xi,\tilde\epsilon^L_i)$ in different volumes, ξ, being the range of $\tilde\epsilon^L_i$, as shown in Fig. 1(b).

The particle density, $\tilde\rho_\xi = \sum_i\tilde\sigma_{\xi i}n_{\xi i}\delta\tilde\epsilon_i$, is represented in Fig. 1(d). Due to the symmetry of the problem, we have pair-wise identical populations, $n_{\xi,i} = n_{N_{\mathbf{r}}-\xi+1,i}$ and $\tilde\rho_\xi = \tilde\rho_{N_{\mathbf{r}}-\xi+1}$, for any ξ and i. The largest deviations in the particle density occurs for the extremal species, placed at both ends of the 1D box as an effect of repulsive interactions.

We observe from Eqs. (4), (6) and from Fig. 1 that the range of $\tilde\epsilon_i$ is $[0,\infty)$ in any elementary volume, ξ. This property of the FES representation allows for particle transport along the system at any energy, ϵ or $\tilde\epsilon$. On the other hand, the range of $\tilde\epsilon^L_{\mathbf{r}_\xi}$ is $[\tilde\epsilon^L_{\xi,\rm min},\infty)$, with $\tilde\epsilon^L_{\xi,\rm min} > 0$ for any ξ. The presence of $\tilde\epsilon^L_{\xi,\rm min}$ creates a non-constant energy gap (Fig. 1b) for the quasi-particle populations and the transport of low energy quasiparticles from one end of the system to the other may occur only through tunneling or higher order processes which involve rearrangements of quasiparticles above the energy gap. In the FES description these rearrangements are taken care of implicitly by the formalism.

Conclusions. – We have formulated an approach by which a system of particles with general particle-particle interaction, $V(|\mathbf{r}-\mathbf{r}'|)$, in an s-dimensional space and external potential $V_{\rm ext}(\mathbf{r})$ is described as an ideal gas of fractional exclusion statistics (FES). We have given the equations for the calculation of the FES parameters and equilibrium populations.

The FES approach have been compared with the Landau's Fermi liquid formalism and we have shown that although there are differences in the definitions of certain quantities like the quasiparticle energies, the physical results are the same. The main difference between the two formalisms is that in the FES approach the quasiparticle energies are independent of the populations of other quasiparticle states and therefore the FES gas is "ideal", with the total energy of the gas being equal to the sum of the quasiparticle energies.

We have exemplified our procedure on a one-dimensional system of fermions with repulsive Coulomb interaction for which we calculated the main microscopic parameters, like the quasiparticle energies, quasiparticle density of states and energy levels populations. For each of these quantities we discussed the similarities and differences between the FES and the FLT approaches.

One practical consequence that appear from our calculations is that the solution of the FES integral equations may eventually be calculated easier by solving self-consistently the FLT equations for population and quasiparticle energies, (3) and (4).

Another consequence is that while in the FLT formulation the quasiparticle energies may form an energy gap at the lowest end of the spectrum due to the particle-particle interaction, in the FES description such an energy gap does not exist. For this reason the low energy particle tunneling in FLT may be described as higher order processes, involving rearrangements of particles on the higher energy levels, wheres in the FES these rearrangements are taken into account directly by the manifestation of the exclusion statistics.

* * *

Enlightening discussions with Francesca Gulminelli and her hospitality during the visit of DVA at the LPC Caen are gratefully acknowledged. The work was supported by the Romanian National Authority for Scientific Research CNCS-UEFISCDI projects PN-II-ID-PCE-2011-3-0960 and PN09370102/2009. The travel support from the Romania-JINR Dubna collaboration project Titeica-Markov are gratefully acknowledged.

REFERENCES

[1] F. D. M. Haldane. *Phys. Rev. Lett.*, 67:937, 1991.
[2] R. Ramanathan. *Phys. Rev. D*, 45:4706, 1992.
[3] A. D. de Veigy and S. Ouvry. *Phys. Rev. Lett.*, 72:600, 1994.
[4] Y.-S. Wu. *Phys. Rev. Lett.*, 73:922, 1994.
[5] S. B. Isakov. *Phys. Rev. Lett.*, 73(16):2150, 1994.
[6] M. V. N. Murthy and R. Shankar. *Phys. Rev. Lett.*, 73:3331, 1994.
[7] M. V. N. Murthy and R. Shankar. Exclusion statistics: From pauli to haldane. Report of The Institute of Mathematical Sciences, Chennai, India, 2013.
[8] S.B. Isakov and S. Viefers. *Int. J. Mod. Phys. A*, 12:1895, 1997.
[9] D. Bernard and Y. S. Wu. In M. L. Ge and Y. S. Wu, editors, *New Developments on Integrable Systems and Long-Ranged Interaction Models*, page 10. World Scientific, Singapore, 1995. cond-mat/9404025.
[10] G. G. Potter, G Müller, and M Karbach. *Phys. Rev. E*, 75:61120, 2007.
[11] G. G. Potter, G Müller, and M Karbach. *Phys. Rev. E*, 76:61112, 2007.
[12] Dan Liu, Ping Lu, Gerhard Müller, and Michael Karbach. *Phys. Rev. E*, 84:021136, 2011.
[13] Dan Liu, Jared Vanasse, Gerhard Müller, and Michael Karbach. *Phys. Rev. E*, 85:011144, 2012.
[14] B. Sutherland. *Phys. Rev. B*, 56:4422, 1997.
[15] K. Iguchi and B. Sutherland. *Phys. Rev. Lett.*, 85:2781, 2000.
[16] D. V. Anghel. *J. Phys. A: Math. Theor.*, 40:F1013, 2007. arXiv:0710.0724.
[17] D. V. Anghel. *Phys. Lett. A*, 372:5745, 2008. arXiv:0710.0728.
[18] D. V. Anghel. *EPL*, 90:10006, 2010. arXiv:0909.0030.
[19] D. Sen and R. K. Bhaduri. *Phys. Rev. Lett.*, 74:3912, 1995.
[20] R. K. Bhaduri, S. M. Reimann, S. Viefers, A. G. Choudhury, and M. K. Srivastava. *J. Phys. B*, 33:3895–3903, 2000.
[21] T. H. Hansson, J. M. Leinaas, and S. Viefers. *Phys. Rev. Lett.*, 86:2930–2933, 2001.
[22] B. Mirza and H. Mohammadzadeh. *Phys. Rev. E*, 78:021127, 2008.
[23] Behrouz Mirza and Hosein Mohammadzadeh. *Phys. Rev. E*, 80:011132, 2009.
[24] B. Mirza and H. Mohammadzadeh. Thermodynamic geometry of fractional statistics. *Phys. Rev. E*, 82:031137, 2010.
[25] Wung-Hong Huang. *Phys. Rev. E*, 51:3729, 1995.
[26] Wung-Hong Huang. *Phys. Rev. B*, 53:15842, 1996.
[27] F. Qin and J. Chen. *J. Phys. B: Atomic, Molecular and Optical Physics*, 43:055302, 2010.
[28] Fang Qin and Ji-sheng Chen. Joule-thomson coefficient of ideal anyons within fractional exclusion statistics. *Phys. Rev. E*, 83:021111, Feb 2011.
[29] Fang Qin and Ji-Sheng Chen. Adiabatic sound velocity and compressibility of a trapped d-dimensional ideal anyon gas. *Phys. Lett. A*, 376:1191, 2012.
[30] Qin Fang and Chen Ji-Sheng. *Commun. Theor. Phys.*, 58:573, 2012.
[31] F. M. D. Pellegrino, G. G. N. Angilella, N. H. March, and

R. Pucci. *Phys. Rev. E*, 76:061123, 2007.

[32] S. B. Isakov, D. P. Arovas, J. Myrheimd, and A. P. Polychronakos. *Phys. Lett. A*, 212:299, 1996.

[33] F. C. Auluck and D. S. Kothari. *Proc. Cambridge Philos. Soc.*, 42:272, 1946.

[34] Robert M. May. *Phys. Rev.*, 135:A1515, 1964.

[35] M. Howard Lee. *Phys. Rev. E*, 55:1518, 1997.

[36] D. V. Anghel. *J. Phys. A: Math. Gen.*, 35:7255, 2002.

[37] G. A. Nemnes and D. V. Anghel. *J. Stat. Mech.*, page P09011, 2010.

[38] I. Guastella, L. Bellomonte, and R. M. Sperandeo-Mineo. *J. Stat. Mech.*, 2009:P02021, 2009.

[39] R. K. Bhaduri, M. V. N. Murthy, and M. K. Srivastava. *J. Phys. B: At. Mol. Opt. Phys.*, 2009.

[40] B. P. van Zyl. *J. Phys. A: Math. Theor.*, 45:315302, 2012.

[41] Z. MacDonald and B. P. van Zyl. *J. Phys. A: Math. Theor.*, 46:045001, 2013.

[42] D. V. Anghel. *EPL*, 87:60009, 2009. arXiv:0906.4836.

[43] D. V. Anghel. *Phys. Rev. Lett.*, 104:198901, 2010.

[44] Yong-Shi Wu. *Phys. Rev. Lett.*, 104:198902, 2010.

[45] D. V. Anghel. *Phys. Lett. A*, 376:892, 2012.

[46] D. V. Anghel. arXiv:1204.4064, 2012.

[47] D. V. Anghel. *Physica Scripta*, 2012:014079, 2012.

[48] D. V. Anghel. *J. Phys. Conf. Ser.*, 410:012121, 2013.

[49] G. A. Nemnes and D. V. Anghel. *J. Phys.: Conf. Ser.*, 410:012120, 2013.